

Government of India

DBT



AIST International Laboratory Advanced Biomedicine



lassroom for Advanced & Frontier Education





Series - 005

Date and Time - Nov 17 (Monday), 4 - 5pm

Venue - Central 4 (5F) Room 5105 Speaker - MAHESH K. KAUSHIK

Affiliation - International Institute for Integrative Sleep Medicine (WPI-IIIS), University of Tsukuba, Japan

E-mail: kaushikmk@gmail.com



Title - Prostaglandin D₂: role in seizure suppression and postictal sleep

Epilepsy is a neurological disorder with the occurrence of seizures, which are often accompanied by sleep. Prostaglandin (PG) D₂ is produced by hematopoietic or lipocalin-type PGD synthase (H- or L-PGDS) and involved in the regulation of physiological sleep. Here, we show that H-PGDS, L/H-PGDS or DP₁ receptor (DP₁R) KO mice exhibited more intense pentylenetetrazole (PTZ)-induced seizures in terms of latency of seizure onset, duration of generalized tonic-clonic seizures, and number of seizure spikes. Seizures significantly increased the PGD₂ content of the brain in wild-type mice. This PTZ-induced increase in PGD₂ was attenuated in the brains of L- or H-PGDS KO and abolished in L/H-PGDS KO mice. Postictal non-rapid eye movement sleep was observed in the wild-type and H-PGDS or DP₂R KO, but not in the L-, L/H-PGDS or DP₁R KO, mice. These findings demonstrate that PGD₂ produced by H-PGDS and acting on DP₁R is essential for seizure suppression and that the L-PGDS/PGD₂/DP₁R system regulates sleep that follows seizures.





Series - 005

Date and Time - Nov 17 (Monday), 4 - Spm Serve - Central 4 (SE) Room, 5105

Venue - Central 4 (SF) Room 510 Speaker - MAHESH K, KAUSHIK

Affiliation – international institute for integrative Sleep Medici ne (WPI-IIIS), University of Tsukuba, Japa E-madi: http://doi.org/10.1006/j.com/



Title - Prostaglandin D₂: role in seizure suppression and postictal sleep

spergy is a reutrological associate with the occurrence of securities, which are often accompanies by sisep. Prostagation (PQI) by produced by hematropoietic or ligicality Ppc PGD synthase (Pt or I-PGDS) and involved in the regulation of physiological sieep. Here, we show that H-PGDS or DP, receptor (DP, 8) No mice exhibited more intense perflementations (PIC)-induced selsiures in terms of latency of selsure onset, duration of generalized tonic-loging selsures, and number of selsures spikes. Selsures significantly increased the PGD, content of the brain in widetype mice. This PICI-induced increase in PGD, was attenuated in the brains of L- or H-PGDS XO and abolished in (H-PGDS XD mice. PRotical non-regid eye movement sleep was observed in the wide-type and A-PGDS or DP, R XD, but not in the L-, L/H-PGDS or DP, R XD, mice. These findings demonstrate that PGD₂ produced by H-PGDS and acting on DP,R is essential for selsure suppression and that the L-PGDS/PGD_/DP,R system regulates selp that follows selsures.



Thank you!

Dear Dr Kaushik

Thanks for being DAILAB-CAFE 005 mentor
We all at BRI-Tsukuba, Hanyang Univ-Seoul, IITDelhi and Brawijaya Univ, Indonesia enjoyed your talk.
Wishing you success for the sleep sacrifice you do for
sleep research...
From.DAILAB